

Treating Painful Diabetic Peripheral Neuropathy: An Update

MATTHEW J. SNYDER, DO, and LAWRENCE M. GIBBS, MD, *Saint Louis University Family Medicine Residency, Belleville, Illinois*

TAMMY J. LINDSAY, MD, *Saint Louis University Family Medicine Residency, St. Louis, Missouri*

Painful diabetic peripheral neuropathy occurs in approximately 25% of patients with diabetes mellitus who are treated in the office setting and significantly affects quality of life. It typically causes burning pain, paresthesias, and numbness in a stocking-glove pattern that progresses proximally from the feet and hands. Clinicians should carefully consider the patient's goals and functional status and potential adverse effects of medication when choosing a treatment for painful diabetic peripheral neuropathy. Pregabalin and duloxetine are the only medications approved by the U.S. Food and Drug Administration for treating this disorder. Based on current practice guidelines, these medications, with gabapentin and amitriptyline, should be considered for the initial treatment. Second-line therapy includes opioid-like medications (tramadol and tapentadol), venlafaxine, desvenlafaxine, and topical agents (lidocaine patches and capsaicin cream). Isosorbide dinitrate spray and transcutaneous electrical nerve stimulation may provide relief in some patients and can be considered at any point during therapy. Opioids and selective serotonin reuptake inhibitors are optional third-line medications. Acupuncture, traditional Chinese medicine, alpha lipoic acid, acetyl-L-carnitine, primrose oil, and electromagnetic field application lack high-quality evidence to support their use. (*Am Fam Physician*. 2016;94(3):227-234. Copyright © 2016 American Academy of Family Physicians.)

CME This clinical content conforms to AAFP criteria for continuing medical education (CME). See CME Quiz Questions on page 208.

Author disclosure: No relevant financial affiliations.

► **Patient information:** A handout on this topic, written by the authors of this article, is available at <http://www.aafp.org/afp/2016/0801/p227-s1.html>.

Painful diabetic peripheral neuropathy (DPN) occurs in approximately 30% of patients with diabetes mellitus who are hospitalized and in 25% of patients with diabetes who are treated in the office setting.¹ It develops as a late manifestation of uncontrolled or long-standing diabetes.¹ As many as 12% of patients with painful DPN do not report symptoms, and 39% of patients with the disorder do not receive any treatment.²

Distal symmetric polyneuropathy, which is characterized by burning pain, paresthesias, and numbness that follows a stocking-glove pattern and progresses proximally, occurs in approximately 26% of patients with DPN. Less than 20% of patients with diabetes experience dynamic mechanical allodynia (pain in response to stroking lightly), thermal hyperalgesia (increased sensitivity to pain by thermal stimuli), or pain attacks.

Poorly controlled blood glucose levels, especially greater variation in glucose levels, contribute to the occurrence and severity of painful DPN.³ A 2012 Cochrane review of two randomized controlled trials (RCTs; n = 1,228) found that enhanced

glucose control in patients with type 1 diabetes significantly reduced the risk of developing DPN (risk difference = 1.84%; 95% confidence interval [CI], -1.11 to -2.56). A similar analysis of four RCTs in patients with type 2 diabetes, however, did not show a statistically significant reduction in the rate of DPN with enhanced glucose control.⁴ Cardiovascular risk factors, including age, tobacco use, hypertension, obesity, dyslipidemia, and peripheral artery disease, are also associated with an increased risk of painful DPN.¹

Pharmacologic and nonpharmacologic interventions are available for the treatment of painful DPN. However, there are few high-quality, head-to-head clinical trials comparing these therapeutic approaches, and because the available studies use varying methodologies, it is difficult to know which treatment strategy may be most effective.

Pharmacologic Treatment Options

Only two medications, pregabalin (Lyrica) and duloxetine (Cymbalta), have been approved by the U.S. Food and Drug Administration (FDA) for the treatment of DPN.

Diabetic Peripheral Neuropathy

Guidelines from national organizations such as the American Academy of Neurology, however, recommend the use of a broader range of medications.

Clinicians should consider the patient's age, health-related quality-of-life goals, physical function, and comorbidities, as well as the possible adverse effects of medication use, in the management of pharmacologic therapy. Additionally, medication dosages and the duration of therapy should be regulated and titrated based on regular patient feedback regarding pain relief, improved function, and adverse effects.¹

Figure 1 presents a treatment algorithm for painful DPN based on available evidence.^{1-3,5-8} Selected medications and their dosages, costs, and numbers needed to treat (NNTs) are presented in Table 1.^{1,7,9-14} Table 2 lists common adverse effects of medications used to treat painful DPN.^{3,12,13,15} Common drug interactions are described in Table 3.¹⁶


ANTICONVULSANTS

Anticonvulsants, including topiramate (Topamax), valproate (Depacon), oxcarbazepine (Trileptal), lamotrigine (Lamictal), and lacosamide (Vimpat), as well as the newer calcium channel alpha-2-delta ligand class anticonvulsants (gabapentin [Neurontin] and pregabalin), have been studied for the treatment of painful DPN. Based on a recent meta-analysis, both the American Academy of Neurology and Toronto guidelines recommend pregabalin as the first-line medication for painful DPN, with gabapentin as the first-line alternative.³

Pregabalin provides consistent pain relief based on a dose-dependent response compared with placebo. A meta-analysis of seven RCTs (n = 1,596) comparing pregabalin with placebo showed that dosages of 300 mg daily and 600 mg daily (or divided twice daily) resulted in greater than 50% pain reduction in 39% and 47% of patients, respectively, vs. 22% with placebo. Onset of action occurred at five days with 300 mg and at four days with 600 mg. Dosing at 150 mg daily was ineffective.¹⁻³

Common adverse reactions to pregabalin (Table 2^{3,12,13,15}) produced discontinuation rates of 9.8% (300 mg) and 14.6% (600 mg).¹

Management of Painful Diabetic Peripheral Neuropathy



May consider adding at any time:

Isosorbide dinitrate spray (30 mg applied to the bottom of feet before bedtime)
Transcutaneous electrical nerve stimulation

ER = extended release; SNRI = serotonin-norepinephrine reuptake inhibitor; SSRI = selective serotonin reuptake inhibitor; TCA = tricyclic antidepressant.

*—Other causes of neuropathy include malignancy, alcohol use, human immunodeficiency virus infection, isoniazid use, and chemotherapy.

†—Pregabalin should be used cautiously in patients with heart failure and in persons concomitantly using thiazolidinediones, because cases of decompensated heart failure with pregabalin use have been reported.¹ In addition, renal insufficiency (creatinine clearance ≤ 30 mL per minute per 1.73 m² [0.50 mL per second per m²]) requires dose adjustment.³

‡—Amitriptyline is the only TCA with good evidence of benefit for treating diabetic neuropathy. TCAs should also be used cautiously and should generally be avoided in older adults because of adverse anticholinergic effects.^{3,6} TCAs are contraindicated in patients with cardiovascular disease (including unstable angina, recent myocardial infarction, heart failure, and abnormal cardiac conduction) and glaucoma.³ Twelve-lead electrocardiography should be performed in all patients older than 40 years to assess for QT prolongation because TCAs may increase the risk of torsades de pointes.

§—May consider combination therapy with pregabalin and gabapentin^{2,5}; however, clinicians should be cautious to avoid serotonin syndrome.

||—Tramadol lowers the seizure threshold and should be avoided in patients with epilepsy or in those at risk of seizures.⁷

¶—Opioids should be used with caution because long-term use is associated with numerous adverse effects and increased morbidity and mortality. If opioids are used, oxycodone is commonly prescribed, but alternatives include methadone, levorphanol, and morphine.

Figure 1. Algorithm for the management of painful diabetic peripheral neuropathy.

Information from references 1 through 3, and 5 through 8.

Table 1. Medications for the Treatment of Painful Diabetic Peripheral Neuropathy

Medication*	Initial dosage	Maximum dosage	Cost†	Comments
Anticonvulsants				
Pregabalin (Lyrica)	300 mg divided twice daily	600 mg divided twice daily	NA (\$370)	NNT = 5 to 8, NNH = 9 to 16‡
Gabapentin (Neurontin)	1,200 mg divided three times daily	3,600 mg divided three times daily	\$10 (\$330)	NNT = 3, NNH = 4‡
Tricyclic antidepressants				
Amitriptyline	10 mg daily at bedtime	150 mg daily at bedtime or divided twice daily	\$14 (NA)	NNT = 1 to 3, NNH = 28 (major adverse events), 6 (minor adverse events)‡
Serotonin-norepinephrine reuptake inhibitors				
Duloxetine (Cymbalta)	60 mg daily or divided twice daily	120 mg daily or divided twice daily	\$32 (\$240)	NNT = 5, NNH = 17‡
Venlafaxine	150 mg daily	225 mg daily	\$14 (\$322)	NNT = 3, NNH = 16
Desvenlafaxine (Pristiq)	200 mg daily	400 mg daily	NA (\$560)	NNT = 9, NNH = 6 to 14‡
Opioid-like medications				
Tramadol	50 mg every 4 to 6 hours	100 mg every 6 hours (maximum 400 mg per day)	\$8 (NA)	NNT = 4, NNH = 8
Tapentadol ER (Nucynta ER)	100 mg twice daily	250 mg twice daily	NA (\$520)	NNT = 9 to 10, NNH = 5‡
Topical medications				
Lidocaine 5% patch	1 patch daily	3 patches daily	\$240 (NA)	NNT = 4
Capsaicin 0.075% cream	Four times daily	NA	NA (\$22 per 55-g tube)	NNT = 7
Isosorbide dinitrate spray	30 mg at bedtime applied to bottom of feet	NA	NA	NA
Selective serotonin reuptake inhibitors				
Citalopram (Celexa)	10 mg daily	40 mg daily	\$6 (\$215)	NNT = 7
Paroxetine (Paxil)	10 mg daily	40 mg daily	\$9 (\$170)	NNT = 7
Escitalopram (Lexapro)	10 mg daily	20 mg daily	\$9 (\$250)	NNT = 7
Opioids				
Oxycodone controlled release§	10 mg twice daily	40 mg twice daily	\$175 (NA)	NNT = 6, NNH = 13

ER = extended release; NA = not available; NNH = number needed to harm; NNT = number needed to treat.

*—Listed in order of category and recommended use.

†—Estimated retail price of one month's treatment based on lowest dosage provided. Cost for generic listed first; brand name in parentheses, when available. Information obtained from <https://www.familywise.org/drug-price-look-up-tool/> and www.amazon.com (accessed November 21, 2015).

‡—Varies in dose-dependent fashion.

§—If opioids are used, oxycodone is commonly prescribed, but other agents, including methadone, levorphanol, and morphine, may be considered. Most studies of opioids for treating neuropathic pain are short-term studies; benefit from long-term use is less certain.

Information from references 1, 7, and 9 through 14.

Pregabalin had similar effectiveness to amitriptyline in relieving painful DPN with fewer adverse events (18% vs. 43%, $P < .0001$).¹ Health-related quality-of-life measures including social functioning, mental health, bodily pain, and vitality all improved, and sleep interference decreased with pregabalin administration.^{2,3}

Gabapentin at dosages of 1,200 mg daily or greater is more effective than placebo and has been shown to have an effect similar to pregabalin (8% to 13% reduction in

symptoms) based on nine RCTs ($n = 1,604$). Compared with amitriptyline, similar effectiveness was noted in at least five RCTs.

Pregabalin is preferred to gabapentin as a first-line medication because of the availability of higher-quality studies on its effects, more predictable pharmacokinetics, shorter titration periods, the option for twice-daily dosing, and no dosing adjustment requirement in patients with renal impairment. Nonetheless, gabapentin has few

Table 2. Adverse Effects of Medications Commonly Used to Treat Painful Diabetic Peripheral Neuropathy

Medication	Adverse effect	Patients experiencing adverse effect (%)
Pregabalin (Lyrica)	Somnolence	5 to 40
	Dizziness	5 to 38
	Peripheral edema	4 to 17
	Headache	2 to 13
	Weight gain > 7%	5
Gabapentin (Neurontin)	Somnolence	22 to 48
	Dizziness	22 to 28
	Weight gain > 7%	5
Amitriptyline	Dry mouth	2 to 89
	Insomnia	35
	Fatigue	11 to 34
	Headache	11 to 21
	Dizziness	5 to 16
Serotonin-norepinephrine reuptake inhibitors	Nausea	10 to 32
	Somnolence	8 to 28
	Dizziness	6 to 25
	Constipation	7 to 19
	Dyspepsia	9 to 18
Tramadol	Constipation	30
	Nausea	30
	Drowsiness	30
Tapentadol ER (Nucynta ER)	Nausea	27
	Anxiety	18
	Diarrhea	16
	Dizziness	15
	Vomiting	13
	Constipation	12
	Restlessness	11
	Headache	10
Lidocaine 5% patch	No adverse effect significantly different from placebo	—
Capsaicin 0.075% cream	Burning at application site	54 to 63
Opioids	Constipation	34
	Drowsiness	29
	Nausea	27
	Dizziness	22
	Hyperalgesia	< 20
	Vomiting	12

ER = extended release.

Information from references 3, 12, 13, and 15.

drug interactions and is the less expensive generic option.^{1,3} Compared with pregabalin, which improves numerous quality-of-life measures, the only quality-of-life measures improved with gabapentin are mental health and vitality.²

Few high-quality studies have investigated the effectiveness of valproate, topiramate, carbamazepine (Tegretol), oxcarbazepine, lamotrigine, and lacosamide. Thus, these agents are not recommended for the treatment of painful DPN.^{2,17}

TRICYCLIC ANTIDEPRESSANTS

Amitriptyline is an effective first-line medication for younger adults.^{1,3} A 2011 systematic review of three RCTs studied the effects of amitriptyline vs. placebo in patients with painful DPN. Amitriptyline had 43% more responders (at least 20% pain reduction) than placebo, reducing pain by 58% to 63% on a five-point pain scale.² Lower-quality evidence from a 2015 Cochrane review of four RCTs including 382 patients with all causes of neuropathic pain showed that amitriptyline use led to a statistically significant benefit (relative risk = 2.0; 95% CI, 1.5 to 2.8; NNT = 5).¹⁸ Health-related quality-of-life outcomes were not studied.

Despite its potential for benefit, amitriptyline is associated with significant adverse anticholinergic effects in up to one-third of patients (*Table 2*^{3,12,13,15}), and its use should be avoided in older adults and monitored with caution in other patients. For other tricyclic antidepressants (i.e., imipramine, desipramine, and nortriptyline [Pamelor]), 11 RCTs provide insufficient high-quality evidence to determine whether they are effective treatments for painful DPN.^{2,19,20}

SEROTONIN-NOREPINEPHRINE REUPTAKE INHIBITORS

Duloxetine is a first-line medication that received FDA approval in 2004 for treating painful DPN.^{1-3,21} A 2014 Cochrane review of eight studies (n = 2,728) showed that duloxetine at 60 mg daily led to at least 50% pain reduction at 12 weeks (NNT = 5; 95% CI, 4 to 7) vs. placebo.⁹ Furthermore, a 2011 multicenter RCT (n = 272) showed duloxetine

Table 3. Common Drug Interactions in the Treatment of Painful Diabetic Peripheral Neuropathy

Medication	Interaction
Pregabalin (Lyrica) and gabapentin (Neurontin)	May increase toxic effects of CNS depressants,* SSRIs, thiazolidinediones Toxic effects may be increased by CNS depressants,* AcCHS inhibitors, hydroxyzine, and magnesium sulfate
Amitriptyline	May increase toxic effects of CNS depressants,* QTc-prolonging agents,† SSRIs, St. John's wort, sulfonyleureas, tramadol, and warfarin (Coumadin) Effectiveness can be decreased by St. John's wort, AcCHS inhibitors, and carbamazepine (Tegretol) Toxic effects can be increased by CNS depressants,* QTc-prolonging agents,† SSRIs, St. John's wort, antiemetics (e.g., 5-HT ₃ antagonists, ondansetron [Zofran]), antipsychotic agents, and bupropion (Wellbutrin) Absolutely contraindicated with MAO inhibitors
Duloxetine (Cymbalta)	May increase toxic effects of CNS depressants,* antiplatelet agents, alpha and beta agonists, anticoagulants (e.g., warfarin), antipsychotic agents Absolutely contraindicated with MAO inhibitors May decrease effects of TCAs and linezolid (Zyvox) Toxic effects may be increased by CNS depressants,* alcohol, antiemetics (e.g., 5-HT ₃ antagonists, ondansetron), antipsychotic agents, metoclopramide (Reglan) Effectiveness may be decreased by CYP1A2 inducers‡
Venlafaxine	May increase toxic effects of CNS depressants* and SSRIs May decrease effectiveness of alpha-2 agonists Absolutely contraindicated with MAO inhibitors and linezolid Toxic effects may be increased by CNS depressants,* CYP2D6 inhibitors,§ CYP3A4 inhibitors, and SSRIs
Tramadol	May increase toxic effects of CNS depressants* and SSRIs Toxic effects may be increased by CNS depressants,* SSRIs, and TCAs
Tapentadol ER (Nucynta ER)	Absolutely contraindicated with MAO inhibitors and mixed agonist and antagonist opioids May increase toxic effects of CNS depressants,* diuretics, and desmopressin Toxic effects may be increased by alcohol, amphetamines, anticholinergic agents, and magnesium sulfate
Opioids (oxycodone controlled release)	May increase toxic effects of CNS depressants,* MAO inhibitors, and SSRIs Effectiveness of medication can be increased by CYP3A4 inducers¶ Toxic effects may be increased by CNS depressants*

AcCHS = acetylcholinesterase; CNS = central nervous system; CYP = cytochrome P450; ER = extended release; MAO = monoamine oxidase; SSRI = selective serotonin reuptake inhibitor; TCA = tricyclic antidepressant.

*—Opioids, zolpidem (Ambien), pregabalin, gabapentin, TCAs, and tramadol.

†—Amiodarone, azithromycin (Zithromax), clarithromycin (Biaxin), fluoxetine (Prozac), haloperidol, risperidone (Risperdal), and sotalol (Betapace).

‡—Carbamazepine, phenobarbital, and rifampin.

§—Amiodarone, celecoxib (Celebrex), cimetidine (Tagamet), clomipramine (Anafranil), desipramine, duloxetine, haloperidol, imipramine, isoniazid, ketoconazole, lidocaine, methadone, pioglitazone (Actos), and sertraline (Zoloft).

||—Amiodarone, cimetidine, clotrimazole, cyclosporine (Sandimmune), desipramine, diltiazem, efavirenz (Sustiva), erythromycin, fluconazole (Diflucan), grapefruit juice, haloperidol, metronidazole (Flagyl), sertraline, tetracycline, and verapamil.

¶—Carbamazepine, dexamethasone, efavirenz, fosphenytoin (Cerebyx), phenytoin (Dilantin), and St. John's wort.

Information from reference 16.

at 60 mg daily to be noninferior to pregabalin at 300 mg daily in patients with an inadequate pain response to treatment with gabapentin over a 12-week period.²² Duloxetine has also been shown to be effective and tolerable in the management of painful DPN in older adults.²³

Other serotonin-norepinephrine reuptake inhibitors (SNRIs) that are used for the treatment of painful DPN include venlafaxine and desvenlafaxine (Pristiq). Although they are considered second-line medications, these SNRIs are often better tolerated and have fewer

drug interactions than amitriptyline, one of the first-line medications.

A 2004 trial showed that higher doses of venlafaxine (150 to 225 mg) led to greater improvements in pain scores than placebo.²⁴ A 2007 Cochrane review of three studies of venlafaxine for general neuropathic pain showed an NNT of 3 (95% CI, 2.2 to 5.1).¹⁰

A 2014 RCT (n = 412) showed desvenlafaxine at dosages of 200 and 400 mg daily to be more effective than placebo in relieving pain associated with DPN, based on a decrease

SORT: KEY RECOMMENDATIONS FOR PRACTICE

<i>Clinical recommendation</i>	<i>Evidence rating</i>	<i>References</i>
Pregabalin (Lyrica), gabapentin (Neurontin), amitriptyline (except in older adults), or duloxetine (Cymbalta) should be used as first-line treatment for painful diabetic peripheral neuropathy.	A	1-3, 8, 9, 22
The serotonin-norepinephrine reuptake inhibitors venlafaxine and desvenlafaxine (Pristiq) and opioid-like medications (tramadol and extended-release tapentadol [Nucynta ER]) may be considered if first-line medications do not provide adequate pain relief.	A	1-3, 6, 8-11
Lidocaine 5% patch, capsaicin 0.075% cream, isosorbide dinitrate spray, and transcutaneous electrical nerve stimulation can be added to therapy for painful diabetic peripheral neuropathy.	B	2, 8, 15, 27, 40

A = consistent, good-quality patient-oriented evidence; B = inconsistent or limited-quality patient-oriented evidence; C = consensus, disease-oriented evidence, usual practice, expert opinion, or case series. For information about the SORT evidence rating system, go to <http://www.aafp.org/afpsort>.

of approximately one point on a 10-point pain rating scale, as well as in improving activity impairment at all doses.¹¹

OPIOID-LIKE MEDICATIONS

Tramadol, a synthetic, centrally-acting opioid-like medication, has also been studied for the treatment of painful DPN. A 2006 Cochrane review of four trials comparing tramadol with placebo showed a significant reduction in neuropathic pain with tramadol at daily dosages of 100 to 400 mg.⁷ A subsequent meta-analysis of three of these trials (n = 303) found an NNT of 4 (95% CI, 2.8 to 6.3) for at least 50% pain relief with tramadol vs. placebo.⁷

Extended-release tapentadol (Nucynta ER), another opioid-like medication, received FDA approval in 2012 for use in the treatment of general neuropathic pain (i.e., not specifically for painful DPN).²⁵ In 2011, a 12-week study (n = 395) demonstrated statistically significant pain relief in patients who were treated with tapentadol (30% pain reduction = 65%; 50% pain reduction = 34.9%; least-squares mean difference of -1.3; 95% CI, -1.70 to -0.92; *P* < .001) vs. placebo.¹³ These positive findings were largely echoed by a later 12-week study comparing tapentadol at dosages of 100 to 250 mg twice daily vs. placebo.²⁶

Topical Medications

Isosorbide dinitrate spray, lidocaine 5% patch or plaster, and capsaicin 0.075% cream should also be considered for the treatment of painful DPN.²

ISOSORBIDE DINITRATE SPRAY

Isosorbide dinitrate is a vasodilator and smooth-muscle relaxant that produced an 18% pain score reduction when sprayed on the skin of areas affected by painful DPN in one small RCT of 22 patients.^{2,27} It can be introduced at any time as an adjunctive treatment.

LIDOCAINE

Lidocaine 5% patch or medicated plaster treatment resulted in comparable pain relief (30% to 50% response) to that of pregabalin (66.7% vs. 69.1%; one

RCT, n = 204). Greater improvement in quality-of-life scores and fewer adverse effects occurred in the lidocaine group than in the pregabalin group.²⁸

CAPSAICIN

Treatment with capsaicin 0.075% cream applied to the painful area four times daily resulted in significant pain relief (50% pain reduction; odds ratio [OR] = 2.37; 95% CI, 1.32 to 4.26) but also adverse events that led to withdrawal from the study (OR = 4.02; 95% CI, 1.45 to 11.06).²⁷ A 2014 systematic review of three RCTs revealed the superiority of capsaicin 0.075% to placebo (standardized mean deviation [SMD] = -0.91; 95% credible interval [CrI], -1.18 to -0.08, large), but not to tricyclic antidepressants (SMD = -0.13; 95% CrI, -1.03 to 0.74, small) or SNRIs (SMD = 0.45; 95% CrI, -0.49 to 1.36, small).¹⁵ The primary adverse effect is burning at the application site, which improves with longer duration of treatment.

Other concentration levels and combination treatments have also been studied. Notably, combining capsaicin 0.025% with doxepin 3.3% led to decreased skin irritation compared with capsaicin alone (one RCT, n = 200, 61% vs. 81% reduction in skin irritation).²⁹

Alternative Medications

SELECTIVE SEROTONIN REUPTAKE INHIBITORS

Selective serotonin reuptake inhibitors are considered third-line medications for treatment of painful DPN because of a lack of high-quality studies supporting their benefit.⁵ Citalopram (Celexa), paroxetine (Paxil), and escitalopram (Lexapro) appear to be mildly effective for treating painful DPN, whereas fluoxetine (Prozac) does not. Combined data from four small studies reveal an NNT of 7 for pain reduction with this class of medications.³⁰

OPIOIDS

Despite concerns about dependency and lower-quality evidence for the benefit of chronic opioid therapy for nonmalignant pain,¹² guidelines suggest a possible

benefit for patients with painful DPN.^{2,5,31} However, monotherapy with opioids is not a first-line treatment and should be reserved for patients who do not achieve pain relief goals and functional improvement with other therapies.^{2,5} Notably, most studies on the use of opioids for neuropathic pain are short-term studies.¹⁴ Chronic opioid use leads to tolerance and hyperalgesia.³²

A 2013 Cochrane review of 14 RCTs (n = 845, 12-week duration or less) analyzed the use of opioids for general neuropathic pain. The review evaluated the effectiveness of methadone, levorphanol, morphine, and controlled-release oxycodone and found that opioids demonstrated superior pain relief compared with placebo (greater than 33% pain relief = 57% vs. 34%; NNT = 4; 95% CI, 2.7 to 7.7; greater than 50% pain relief = 47% vs. 30%; NNT = 6; 95% CI, 3.0 to 50.0).¹² Treatment with opioids did not, however, result in improvement in many aspects of emotional or physical functioning, as measured by various validated questionnaires. In contrast, a 2014 Cochrane review assessing the use of controlled-release oxycodone (three studies, n = 204) in the treatment of painful DPN found insufficient evidence to support its role in therapy.³³

Other Treatments

ACUPUNCTURE

Acupuncture, which has been the focus of 75 low-quality RCTs; traditional Chinese medicine, which typically includes acupuncture and herbal medicine; and electromagnetic field application are not currently recommended for the treatment of painful DPN because large, high-quality RCTs are lacking.^{2,34-36}

ALPHA LIPOIC ACID

Alpha lipoic acid is an antioxidant substance that has been studied for treatment of neuropathic pain. Based on a 2012 meta-analysis (n = 448), alpha lipoic acid (600 mg) administered intravenously daily for three weeks led to a decrease in reported neuropathic pain for up to one year (standardized mean symptom score reduction = -2.81; 95% CI, -4.16 to -1.46, *P* = .0001). However, studies investigating long-term clinical improvement are lacking. Oral administration did not produce a clinically significant symptom score reduction.^{1,37}

ACETYL-L-CARNITINE

Acetyl-L-carnitine, based on a 2008 meta-analysis of two RCTs (n = 1,679), and primrose oil, based on three small RCTs, demonstrated an improvement in nerve conduction but a lack of clinically significant long-term improvement in symptom scores.^{38,39}

TRANSCUTANEOUS ELECTRICAL NERVE STIMULATION

The results of several small studies suggest that transcutaneous electrical nerve stimulation (TENS) should be considered as a treatment for painful DPN.⁴⁰ An RCT involving 31 participants showed that 15 of 18 (83%) patients receiving TENS had improved pain scores (reduction from 3.17 to 1.44 on a 5-point scale) vs. five of 13 (38%) patients receiving sham treatments (reduction from 2.98 to 2.38).⁴¹ In another small RCT (n = 19), pain scores were reduced by 27% with six weeks of TENS therapy.⁴² TENS can be started at any time during therapy as an adjunct to other treatments.

Data Sources: The primary literature search was completed with Essential Evidence Plus and included searches of the Cochrane database, PubMed, and National Guideline Clearinghouse with the terms diabetic neuropathies and diabetic neuropathy. In addition, a PubMed search was completed using the terms diabetic neuropathy pain treatment, anti-convulsants, tricyclic antidepressant, alpha lipoic acid, primrose oil, and alternative medicine. Search date: November 20, 2015.

The opinions and assertions contained herein are the private views of the authors and are not to be construed as official or as reflecting the views of Saint Louis University, the U.S. Air Force Medical Service, or the U.S. Air Force at large.

This review updates a previous article on this topic by Lindsay, Rodgers, Savath, and Hettinger.⁴³

The Authors

MATTHEW J. SNYDER, DO, FFAFP, is associate program director of the Saint Louis University Family Medicine Residency Program in Belleville, Ill., which is affiliated with Scott Air Force Base.

LAWRENCE M. GIBBS, MD, FFAFP, is an assistant clinical professor in the Saint Louis University Family Medicine Residency Program, Belleville.

TAMMY J. LINDSAY, MD, FFAFP, is an associate clinical professor with the Department of Family and Community Medicine at Saint Louis University School of Medicine, St. Louis, Mo.

Address correspondence to Matthew J. Snyder, DO, Saint Louis University, 180 S. Third St., Suite 400, Belleville, IL 622209 (e-mail: mdrsnyder@gmail.com). Reprints are not available from the authors.

REFERENCES

- Spallone V, Lacerenza M, Rossi A, Sicuteri R, Marchettini P. Painful diabetic polyneuropathy: approach to diagnosis and management. *Clin J Pain*. 2012;28(8):726-743.
- Bril V, England J, Franklin GM, et al.; American Academy of Neurology; American Association of Neuromuscular and Electrodiagnostic Medicine; American Academy of Physical Medicine and Rehabilitation. Evidence-based guideline: Treatment of painful diabetic neuropathy [published correction appears in *Neurology*. 2011;77(6):603]. *Neurology*. 2011;76(20):1758-1765.
- Ziegler D, Fonseca V. From guideline to patient: a review of recent recommendations for pharmacotherapy of painful diabetic neuropathy. *J Diabetes Complications*. 2015;29(1):146-156.
- Callaghan BC, Little AA, Feldman EL, Hughes RA. Enhanced glucose control for preventing and treating diabetic neuropathy. *Cochrane Database Syst Rev*. 2012;(6):CD007543.

Diabetic Peripheral Neuropathy

- Moulin D, Boulanger A, Clark AJ, et al.; Canadian Pain Society. Pharmacological management of chronic neuropathic pain: revised consensus statement from the Canadian Pain Society. *Pain Res Manag.* 2014; 19(6):28-335.
- Vieweg WV, Wood MA, Fernandez A, Beatty-Brooks M, Hasnain M, Pandurangi AK. Proarrhythmic risk with antipsychotic and antidepressant drugs: implications in the elderly. *Drugs Aging.* 2009;26(12):997-1012.
- Hollingshead J, Dühmke RM, Cornblath DR. Tramadol for neuropathic pain. *Cochrane Database Syst Rev.* 2006;(3):CD003726.
- Finnerup NB, Attal N, Haroutounian S, et al. Pharmacotherapy for neuropathic pain in adults: a systematic review and meta-analysis. *Lancet Neurol.* 2015;14(2):62-173.
- Lunn MP, Hughes RA, Wiffen PJ. Duloxetine for treating painful neuropathy, chronic pain or fibromyalgia. *Cochrane Database Syst Rev.* 2014; (1):CD007115.
- Saarto T, Wiffen PJ. Antidepressants for neuropathic pain. *Cochrane Database Syst Rev.* 2007;(4):CD005454.
- Allen R, Sharma U, Barlas S. Clinical experience with desvenlafaxine in treatment of pain associated with diabetic peripheral neuropathy. *J Pain Res.* 2014;(7):39-351.
- McNicol ED, Midbari A, Eisenberg E. Opioids for neuropathic pain. *Cochrane Database Syst Rev.* 2013;(8):CD006146.
- Schwartz S, Etropolski M, Shapiro DY, et al. Safety and efficacy of tapentadol ER in patients with painful diabetic peripheral neuropathy: results of a randomized-withdrawal, placebo-controlled trial. *Curr Med Res Opin.* 2011;27(1):151-162.
- Dowell D, Haegerich TM, Chou R. CDC Guideline for prescribing opioids for chronic pain—United States, 2016. *MMWR Recomm Rep.* 2016;65(1):1-49.
- Griebeler ML, Morey-Vargas OL, Brito JP, et al. Pharmacologic interventions for painful diabetic neuropathy: an umbrella systematic review and comparative effectiveness network meta-analysis [published corrections appear in *Ann Intern Med.* 2015;162(8):600, and *Ann Intern Med.* 2015;162(10):739]. *Ann Intern Med.* 2014;161(9):639-649.
- Lexicomp. online.lexi.com [subscription required]. Accessed March 5, 2015.
- Wiffen PJ, Derry S, Moore RA, Kalso EA. Carbamazepine for chronic neuropathic pain and fibromyalgia in adults. *Cochrane Database Syst Rev.* 2014;(4):CD005451.
- Moore RA, Derry S, Aldington D, Cole P, Wiffen PJ. Amitriptyline for neuropathic pain in adults. *Cochrane Database Syst Rev.* 2015;(7):CD008242.
- Hearn L, Derry S, Phillips T, Moore RA, Wiffen PJ. Imipramine for neuropathic pain in adults. *Cochrane Database Syst Rev.* 2014;(5):CD010769.
- Hearn L, Moore RA, Derry S, Wiffen PJ, Phillips T. Desipramine for neuropathic pain in adults. *Cochrane Database Syst Rev.* 2014;(9):CD011003.
- Bymaster FP, Dreshfield-Ahmad LJ, Threlkeld PG, et al. Comparative affinity of duloxetine and venlafaxine for serotonin and norepinephrine transporters in vitro and in vivo, human serotonin receptor subtypes, and other neuronal receptors. *Neuropsychopharmacology.* 2001; 25(6):871-880.
- Tanenberg RJ, Irving GA, Risser RC, et al. Duloxetine, pregabalin, and duloxetine plus gabapentin for diabetic peripheral neuropathic pain management in patients with inadequate pain response to gabapentin: an open-label, randomized, noninferiority comparison. *Mayo Clin Proc.* 2011;86(7):615-626.
- Skljarevski V, Frakes EP, Sagman D, Lipsius S, Heinloth AN, Dueñas-Tentori HJ. Review of efficacy and safety of duloxetine 40 to 60mg once daily in patients with diabetic peripheral neuropathic pain. *Pain Res Treat.* 2012;2012:898347.
- Rowbotham MC, Goli V, Kunz NR, Lei D. Venlafaxine extended release in the treatment of painful diabetic neuropathy: a double-blind, placebo-controlled study [published correction appears in *Pain.* 2005;113(1-2):248]. *Pain.* 2004;110(3):697-706.
- Tzschentke TM, Christoph T, Kögel B, et al. (-)-(1R,2R)-3-(3-dimethylamino-1-ethyl-2-methyl-propyl)-phenol hydrochloride (tapentadol HCl): a novel mu-opioid receptor agonist/norepinephrine reuptake inhibitor with broad-spectrum analgesic properties. *J Pharmacol Exp Ther.* 2007;323(1):265-276.
- Vinik AI, Shapiro DY, Rauschkolb C, et al. A randomized withdrawal, placebo-controlled study evaluating the efficacy and tolerability of tapentadol extended release in patients with chronic painful diabetic peripheral neuropathy. *Diabetes Care.* 2014;37(8):2302-2309.
- Wong MC, Chung JW, Wong TK. Effects of treatments for symptoms of painful diabetic neuropathy: systematic review. *BMJ.* 2007; 335(7610):87.
- Baron R, Mayoral V, Leijon G, Binder A, Steigerwald I, Serpell M. 5% lidocaine medicated plaster versus pregabalin in post-herpetic neuralgia and diabetic polyneuropathy: an open-label, non-inferiority two-stage RCT study. *Curr Med Res Opin.* 2009;25(7):1663-1676.
- Argoff CE. Topical analgesics in the management of acute and chronic pain. *Mayo Clin Proc.* 2013;88(2):195-205.
- Finnerup NB, Sindrup SH, Jensen TS. The evidence for pharmacological treatment of neuropathic pain. *Pain.* 2010;150(3):573-581.
- Chou R, Fanciullo GJ, Fine PG, et al.; American Pain Society-American Academy of Pain Medicine Opioids Guidelines Panel. Clinical guidelines for the use of chronic opioid therapy in chronic noncancer pain. *J Pain.* 2009;10(2):113-130.
- Chang G, Chen L, Mao J. Opioid tolerance and hyperalgesia. *Med Clin North Am.* 2007;91(2):199-211.
- Gaskell H, Moore RA, Derry S, Stannard C. Oxycodone for neuropathic pain and fibromyalgia in adults. *Cochrane Database Syst Rev.* 2014;(6):CD010692.
- Weintraub MI, Herrmann DN, Smith AG, Backonja MM, Cole SP. Pulsed electromagnetic fields to reduce diabetic neuropathic pain and stimulate neuronal repair: a randomized controlled trial. *Arch Phys Med Rehabil.* 2009;90(7):1102-1109.
- Galuppo M, Giacoppo S, Bramanti P, Mazzon E. Use of natural compounds in the management of diabetic peripheral neuropathy. *Molecules.* 2014;19(3):877-2895.
- Chen W, Yang GY, Liu B, Manheimer E, Liu JP. Manual acupuncture for treatment of diabetic peripheral neuropathy: a systematic review of randomized controlled trials [published correction appears in *PLoS One.* 2014;9(3):e91110]. *PLoS One.* 2013;8(9):e73764.
- Mijnhout GS, Kollen BJ, Alkhalaf A, Kleefstra N, Bilo HJ. Alpha lipoic acid for symptomatic peripheral neuropathy in patients with diabetes: a meta-analysis of randomized controlled trials. *Int J Endocrinol.* 2012; 2012:456279.
- Evans JD, Jacobs TF, Evans EW. Role of acetyl-L-carnitine in the treatment of diabetic peripheral neuropathy. *Ann Pharmacother.* 2008; 42(11):686-1691.
- Halat KM, Dennehy CE. Botanicals and dietary supplements in diabetic peripheral neuropathy. *J Am Board Fam Pract.* 2003;16(1):47-57.
- Dubinsky RM, Miyasaki J. Assessment: efficacy of transcutaneous electrical nerve stimulation in the treatment of pain in neurologic disorders (an evidence-based review): report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *Neurology.* 2010;74(2):173-176.
- Kumar D, Marshall HJ. Diabetic peripheral neuropathy: amelioration of pain with transcutaneous electrostimulation. *Diabetes Care.* 1997;20(11):1702-1705.
- Forst T, Nguyen M, Forst S, Disselhoff B, Pohlmann T, Pfützner A. Impact of low frequency transcutaneous electrical nerve stimulation on symptomatic diabetic neuropathy using the new Salutaris device. *Diabetes Nutr Metab.* 2004;17(3):163-168.
- Lindsay TJ, Rodgers BC, Savath V, Hettinger K. Treating diabetic peripheral neuropathic pain. *Am Fam Physician.* 2010;82(2):151-158.